

Prävalenz und Inzidenz M31

insbesondere Wegener-Granulomatose M31.3

Folien von
Frau PD Dr. med. E. Reinhold-Keller
Hamburg 08.03.2008

Prävalenz der WG in- und außerhalb von Europa: Neue Ergebnisse aus Schweden

	NZL	US	N	F	UK	D	S
WG	152	30	49-95	24	63	58	160
MPA	58					9	94
Studienperiode	'99-03	79-88	93-98	2000	1997	1995	2003

**In Schweden wurden die bislang höchsten
Prävalenzdaten für die WG ermittelt**

Inzidenz der WG in- und außerhalb von Europa:

Neue Ergebnisse aus UK und D

	S	UK	UK	E	N	US	D	Peru
Periode	75-01	88-98	00-05	88-01	84-98	90-99	98-06	90-04
Stud.Typ	popul.	hosp.	popul.	hosp.	popul.	popul.	popul.	popul.
Population, n	8 Mio.	413.000	2.8 Mio	208.271	464.000		2.8 Mio.	930.306

Inzidenz ANCA-assoziiierter Vaskulitiden (Neuerkrankungen/1 Mio/J.)

WG	7.8	10.6	8.0	2.9	8.0	8.3	9.0	0.50
CSS		3	2.5	1.3			1	
MPA		8.4		7.91			3	4.01

- Cotch MF, Hoffman GS, Yerg DE, Kaufman GI, Targonski P, Kaslow RA. The epidemiology of Wegener`s granulomatosis. *Arthritis Rheum* 1996; 39: 87-92
- Dadoniene J, Kirdaite G, Mackiewicz Z Rimkevicius A, Haugeberg G: Incidence fo primary systemic vasculitides in Vilnius : a university hospital population based study. *Ann Rheum Dis* 2005; 64: 335-336
- Gibson A, Stamp LK, Chapman PT, O`Donnell JL: The epidemiology of Wegener`s granulomatosis and microscopic polyangiitis in a Southern hemisphere region. *Rheumatology* 2006; 45: 624-628
- Gonzalez-Gay MA, Garcia-Porrúa C, Guerrero J, Rodriguez-Ledo P, Llorca J: The epidemiology of the primary systemic vasculitides in northwest Spain: implications of the Chapel Hill Concensus Conference Defintions. *Arthritis Rheum* 2003; 49: 388-393
- Herlyn K, Gross WL, Reinhold-Keller E. Stabile Inzidenzraten der primär-systemischen Vaskulitiden in Schleswig-Holstein von 1998-2006. *ZfRh* 2007; 66 (Suppl. 1): 65
- Knight A, Ekbohm A, Brandt L, Askling J. Increasing incidence of Wegener`s granulomatosis in Sweden, 1975-2001. *J Rheumatol* 2006, 33: 2060-3.
- Koldingsnes W, Nossent H: Epidemiology of Wegener`s Granulomatosis in northern Norway. *Arthritis Rheum* 2000; 43:2481-2487
- Lane SE, Scott DG, Heaton A, Watts RA: Primary renal vasculitis in Norfolk : increasing incidence or increasing recognition. *Nephrol Dial Transplant* 2000; 15: 23-27
- Mahr A, Guillevin L, Poissonnet M, Ayme S. Prevalence of polyarteritis nodosa, microscopic polyangiitis, Wegener`s granulomatosis and Churg Strauss syndrome in a Frenche urban multiethnic population in 2000, a capture-recapture estimate. *Arthritis Care & Res* 2004; 51: 92-99
- Mohammad AJ, Jacobsson LTH, Mahr AD, Sturfelt G, Segelmark M. Prevalence of Wegener`s granulomatosis, microscopic polyangiitis, polyarteritis nodosa and Churg Strauss syndrome within a defined population in southern Sweden. *Rheumatology* 2007; 46: 1329-1337.
- Reinhold-Keller E, Herlyn K, Wagner-Bastmeyer R, Gross WL: Stable incidence of primary systemic vasculitides over five years: Results from the German vasculitis register. *Arthritis Rheum* 2005; 53: 93-99
- Reinhold-Keller E, Herlyn K, Wagner-Bastmeyer R, Gutfleisch J, Peter HH, Raspe HH, Gross WL: No differences in the incidence of vasculitides between north and south Germany : first results of the German vasculitis register. *Rheumatology* 2002; 41: 540-549
- Reinhold-Keller E, Zeidler A, Gutfleisch J, Peter HH, Raspe HH, Gross WL. Giant cell arteritis is more prevalent in urban than in rural population: results of an epidemiological study of primary systemic vasculitides in Germany. *Rheumtology* 2000; 39: 1396-1402.
- Watts RA: What is known about the epidemiology of the vasculitides. *Best Pract Res Clin Rheum* 2005; 19:191-207
- Watts R, Al-Taiar A, Skinner J, Scott D, MacGregor A. The incidence of Wegener`s granulomatosis and Churg Strauss syndrome in the UK General Practice Database. *ACR* 2007.
- Watts RA, Gonzalez-Gay MA, Lane SE, Garcia-Porrúa C, Bentham G, Scott DG: Geoepidemiology of systemic vasculitis: comparison of the incidence in two regions of Europa. *Ann Rheum Dis* 2001; 60: 170-172
- Watts R, Lane S, Bentham G, Scott D. Epidemiology of systemic vasculitides. A ten year study in the United Kingdom. *Arthritis Rheum* 2000; 43: 414-419